

Tip:

You can view the presentation in full screen mode and change pages by using the arrow keys or the scroll wheel of the mouse.

To get into full screen mode, choose [View] on the menu and scroll down to [Full Screen] or simply press the key combination [Ctrl]+[L].
To leave the full screen mode, press the [Esc] key.

www.puls-schlag.org

Please distribute this information widely

Pass this information on to:

Medical and complimentary practitioners

MPs and councillors

Planning officers

Local park and forestry authorities

Gardeners, horticultural societies, tree surgeons and nurseries

Local health authorities

Local government ecologists

Local environmental protection and conservation associations

Teachers, school governors and heads of schools

Friends and family

P))) **ULS-SCHLAG**

presents

Tree Damage from Chronic High Frequency Exposure?

Mobile Telecommunications, Radar, Point-to-point transmission systems,
Terrestrial Radio and TV etc.

Timeline Sequence: „The Intelligencetest“

Issued: May 2007



[Dr.-Ing. Dipl.-Phys. Volker Schorpp](#)

© P))) ULS-SCHLAG e.V. Karlsruhe, Germany

www.puls-schlag.org

Intelligence Test

Where is the transmitter?

White Horse Chestnut Tree



Karlsruhe, 10.09.2006

Karlsruhe, 10.09.2006

Why would any pest attack trees just at that side were they are exposed to a HF-Transmitter?

Panorama Shot

HF-Transmitter



Karlsruhe, 08.10.2006

Panorama Shot

HF-Transmitter



2 months later

Karlsruhe, 01.11.2006

Panorama Shot

HF-Transmitter



Exposed



"Protected"
trees



Another 3 weeks later

Comments to the „Intelligence Test“

Already in summer, the large white horse chestnut tree exhibits damage on the side of the transmitter in the form of brown coloured leaves which fall off. The damage grows in the direction of the radiation until the tree is completely bare after a few weeks. The browning of the leaves is not patchy as it would be due to an infestation of horse chestnut leaf-miner (*cameraria ohridella*), but instead it starts at the edge of the leaves and expands towards the leaf stems. The leaves appear to be literally drying up.

The large tree to the right of the horse chestnut (lyme tree) does not share the same fate. Apparently, not every type of HF exposure (HF field configuration) is affecting every tree in the same way.

The trees behind the tall building, which are more protected from HF exposure, exhibit a later and spatially more homogenous wilting behaviour.

Tree Damage from Chronic High Frequency Exposure

More informations and explanations at

www.puls-schlag.org

Please support P)))ULS-SCHLAG

IBAN	DE37 6609 0800 0005 366097
BIC (SWIFT-Code)	GENODE61BBB
Bank	BBBank Karlsruhe

Please support our campaign for life!

www.puls-schlag.org

The End

www.puls-schlag.org