Tip:

You can view the presentation in full screen mode and change pages by using the arrow keys or the scroll wheel of the mouse.

To get into full screen mode, choose [View] on the menu and scroll down to [Full Screen] or simply press the key combination [Ctrl]+[L]. To leave the full screen mode, press the [Esc] key.

Please distribute this information widely

Pass this information on to:

Medical and complimentary practitioners MPs and councillors Planning officers Local park and forestry authorities Gardeners, horticultural societies, tree surgeons and nurseries Local health authorities Local government ecologists Local environmental protection and conservation associations Teachers, school governors and heads of schools Friends and family

P)))ULS-SCHLAG

presents

Tree Damage from Chronic High Frequency Exposure?

Mobile Telecommunications, Radar, Point-to-point Transmission Systems, Terrestrial Radio and TV etc.

Timeline Sequence: "The Three Lime Trees" Issued: May 2007



Dr.-Ing. Dipl.-Phys. Volker Schorpp

© **P**))**ULS-SCHLAG e.V.** Karlsruhe, Germany

The Three Lime Trees Could you think of any better way for the trees to point to the cause for their disease?

Translation from German by Andrea Klein, London



© **P**))**ULS-SCHLAG** Karlsruhe, Germany

















Typically unnatural leaf wilt in the tree top area exposed to high frequency radiation

Spatially irregular exposure of free standing trees occurs almost exclusively in the built-up environment. Therefore, irregular damage patterns as shown in this picture are also almost exclusively found in the built-up environment.



(CP)))**ULS-SCHLAG** Karlsruhe, Germany



Explanatory Model for the Timeline Sequence "The Three Lime Trees"

Mainly due to their position within the built-up environment, the three lime trees are exposed in different ways. They exhibit spatially different damage structures with different damage timelines. The exposure conditions in this case are simple and straightforward for anyone familiar with the propagation of high frequency radiation: The radiation is diffracted (bent downwards) by the roof ridge of the light blue house. Lime tree number 1(left) has the highest position. Its transmitter facing side suffers full height exposure to the diffracted high frequency radiation. The exhibited damage is also transmitter facing, i.e. it originates on the side facing the source of the radiation and expands in the direction of the radiation. The smaller lime tree number 2 has the lowest position and is shielded by the surrounding buildings. The propagation path of the diffracted radiation goes straight past its tree top. This tree does not exhibit any transmitter facing damage and sheds its leaves only in mid-November and with a regular pattern of wilting. Lime tree number 3 is positioned in a way which exposes part of its tree top to the diffracted radiation from the transmitter. This tree exhibits the "typical" spatially inhomogeneous (irregular) damage, expressed by the unnatural premature wilting of the leaves in the tree top. A situation where free standing trees are only partially exposed to high frequency radiation (i.e. only in their tree tops) can usually only be found within the built-up environment. Hence, this particular pattern of damage is also almost exclusively found in free standing deciduous trees within a builtup environment.

V. Schorpp

(CP))**ULS-SCHLAG** Karlsruhe, Germany

Tree Damage from Chronic High Frequency Exposure

More informations and explanations at

Please support **P**)) ULS-SCHLAG

IBANDE37 6609 0800 0005 366097BIC (SWIFT-Code)GENODE61BBBBankBBBank Karlsruhe

Please support our campaign for life!

The End